## COMPSOCRYPTUS JAMIESONI, A NEW ICHNEUMONID FROM CALIFORNIA (HYMENOPTERA)<sup>1</sup>

## Sal Nolfo<sup>2</sup>

ABSTRACT: Compsocryptus jamiesoni, n. sp. is described from Alviso (Santa Clara County), California.

Compsocryptus jamiesoni, n. sp.

Male — Fore wing 11 mm. long: mesoscutum with punctures moderately coarse and strong, separated by an average of 0.5 their diameter.

Ferruginous. Area on the middle of frons and vertex, occiput, apical 0.4 of flagellum, sutural markings of thorax, black; hind tibia infuscate apically; base of third tergite fuscous; wings yellowish with thee transverse dark brown bands.

Female — Fore wing 9.5 mm. to 10 mm. long: temple at midheight about 0.82 as long as eye, moderately convex; ovipositor about 6.1 as long from nodus to apex as it is deep at nodus. Ferruginous. Base of third tergite and narrow areas along some of thoracic sutures black; apical 0.45 of flagellum fuscous; wings brownish yellow with thee transverse dark brown bands; hind tibia infuscate apically.

Holotype —  $\mathfrak{P}$ , Alviso (Santa Clara County), Catifornia, June 23, 1981. Dean Jamieson, in Salt Marsh. (This holotype is on indefinite loan from the Entomology Museum, North Carolina State University, Raleigh, to the United States National Museum, Wash., D.C.). Paratypes —  $2\mathfrak{P}$ ,  $2\mathfrak{P}$ , Alviso, California, from June 23-29, 1981, Dean Jamieson.

This species closely resembles both Compsocryptus aridus and Compsocryptus calipterus. Because the female has sparse hairs on the second tergite, it keys to Compsocryptus aridus in Townes (1962) keys. The female is similar to Compsocryptus aridus but differs in having the body color ferruginous rather than fulvous, dark markings on the wings broader, and a slightly larger average size. Except for the sparse hairs on the abdominal tergite, the male is similar to Compsocryptus calipterus but lacks the dark area at the apex of the hind femur. Compsocryptus aridus are found in deserts. Compsocryptus jamiesoni inhabit salt marsh areas.

## ACKNOWLEDGMENTS

I wish to thank Dean Jamieson for providing the 5 specimens examined. I am also grateful to Dr. Henry Townes, of the American Entomological Institute, whose advice was most helpful.

## REFERENCE

Townes, Henry, 1962, U.S. Nat. Mus. Bull. No. 216, pt. 3, pages 278-306.

ENT. NEWS, 93 (2) 42, March & April 1982

<sup>&</sup>lt;sup>1</sup>Received September 22, 1981

<sup>&</sup>lt;sup>2</sup>1419 Valley Rd., Garner, North Carolina 27529